

***Haemophilus influenzae*, Invasive Cases^b and Incidence Rates^a in Texas by Age Group, 2007-2017**

| Age Group | 2007 | | 2008 | | 2009 | | 2010 | | 2011 | | 2012 | | 2013 | | 2014 | | 2015 | | 2016 | | 2017 | |
|-----------|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|-------|-----|
| | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR | Cases | IR |
| <1 | 3 | 0.8 | 2 | 0.5 | 2 | 0.5 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.2 | 4 | 1.0 | 3 | 0.7 | 26 | 6.2 | 29 | 6.7 |
| 1-4 | 0 | 0.0 | 0 | 0.0 | 1 | 0.1 | 2 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 2 | 0.1 | 1 | 0.1 | 17 | 1.0 | 21 | 1.3 |
| 5-9 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.1 | 0 | 0.0 | 0 | 0.0 | 1 | 0.1 | 0 | 0.0 | 1 | 0.1 | 12 | 0.6 | 4 | 0.2 |
| 10-14 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 4 | 0.2 | 2 | 0.1 |
| 15-19 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 3 | 0.1 | 5 | 0.2 |
| 20-29 | 0 | 0 | 0 | 0.0 | 0 | 0.0 | 3 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0 | 0.0 | 11 | 0.3 | 17 | 0.4 |
| 30-39 | 0 | 0.0 | 1 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 1 | 0.0 | 0 | 0.0 | 12 | 0.3 | 26 | 0.6 |
| 40-49 | 3 | 0.1 | 0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 3 | 0.1 | 16 | 0.4 | 25 | 0.7 |
| 50-59 | 1 | 0.0 | 0 | 0.0 | 2 | 0.1 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 33 | 0.9 | 34 | 1.0 |
| 60 and Up | 7 | 0.2 | 8 | 0.2 | 2 | 0.1 | 3 | 0.1 | 1 | 0.0 | 2 | 0.1 | 1 | 0.0 | 4 | 0.1 | 2 | 0.0 | 182 | 3.8 | 240 | 4.6 |
| Unknown | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - | 0 | - |

^aIncidence rates (IR) calculated per 100,000 population

^b2007-2015 *Haemophilus influenzae* invasive type b cases reported only; all serotypes reported in 2016-2017