

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries
 Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence [†]	95% Confidence Interval for Prevalence	
Central Nervous System					
Anencephaly	2000	101	2.78	2.24 -	3.32
	2001	96	2.63	2.13 -	3.21
	2002	97	2.60	2.11 -	3.18
	2003	98	2.60	2.11 -	3.16
	2004	88	2.31	1.85 -	2.84
	2005	86	2.23	1.78 -	2.75
	2006	91	2.28	1.83 -	2.80
	2007	93	2.28	1.84 -	2.80
	2008	109	2.69	2.18 -	3.19
	2009	124	3.09	2.54 -	3.63
Spina bifida without anencephaly	2000	138	3.80	3.16 -	4.43
	2001	123	3.37	2.77 -	3.96
	2002	113	3.03	2.48 -	3.59
	2003	122	3.23	2.66 -	3.81
	2004	173	4.54	3.86 -	5.21
	2005	141	3.66	3.05 -	4.26
	2006	136	3.41	2.83 -	3.98
	2007	154	3.78	3.18 -	4.38
	2008	143	3.53	2.95 -	4.11
	2009	151	3.76	3.16 -	4.36
Encephalocele	2000	39	1.07	0.76 -	1.47
	2001	35	0.96	0.67 -	1.33
	2002	28	0.75	0.50 -	1.09
	2003	33	0.87	0.60 -	1.23
	2004	23	0.60	0.38 -	0.90
	2005	33	0.86	0.59 -	1.20
	2006	39	0.98	0.69 -	1.34
	2007	44	1.08	0.78 -	1.45
	2008	41	1.01	0.73 -	1.37
	2009	39	0.97	0.69 -	1.33
Holoprosencephaly	2000	38	1.05	0.74 -	1.44
	2001	46	1.26	0.92 -	1.68
	2002	51	1.37	1.02 -	1.80
	2003	45	1.19	0.87 -	1.60
	2004	37	0.97	0.68 -	1.34
	2005	41	1.06	0.76 -	1.44
	2006	46	1.15	0.84 -	1.54
	2007	54	1.33	1.00 -	1.73
	2008	37	0.91	0.64 -	1.26
	2009	31	0.77	0.52 -	1.10

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence	
Hydrocephaly without spina bifida	2000	265	7.29	6.42 -	8.17
	2001	233	6.38	5.56 -	7.20
	2002	251	6.74	5.91 -	7.57
	2003	249	6.60	5.78 -	7.42
	2004	265	6.95	6.11 -	7.78
	2005	234	6.07	5.29 -	6.85
	2006	269	6.74	5.93 -	7.54
	2007	309	7.58	6.74 -	8.43
	2008	297	7.33	6.50 -	8.16
	2009	293	7.30	6.46 -	8.13
Eye and Ear					
Anophthalmia	2000	11	0.30	0.15 -	0.54
	2001	11	0.30	0.15 -	0.54
	2002	11	0.30	0.15 -	0.53
	2003	12	0.32	0.16 -	0.56
	2004	12	0.31	0.16 -	0.55
	2005	17	0.44	0.26 -	0.71
	2006	15	0.38	0.21 -	0.62
	2007	13	0.32	0.17 -	0.55
	2008	9	0.22	0.10 -	0.42
	2009	13	0.32	0.17 -	0.55
Microphthalmia	2000	91	2.50	2.02 -	3.08
	2001	89	2.44	1.96 -	3.00
	2002	105	2.82	2.28 -	3.36
	2003	101	2.68	2.15 -	3.20
	2004	95	2.49	2.02 -	3.04
	2005	100	2.59	2.11 -	3.15
	2006	121	3.03	2.49 -	3.57
	2007	131	3.22	2.66 -	3.77
	2008	121	2.99	2.45 -	3.52
	2009	101	2.51	2.02 -	3.01
Cataract	2000	63	1.73	1.33 -	2.22
	2001	52	1.42	1.06 -	1.87
	2002	75	2.01	1.58 -	2.52
	2003	65	1.72	1.33 -	2.20
	2004	71	1.86	1.45 -	2.35
	2005	77	2.00	1.58 -	2.50
	2006	81	2.03	1.61 -	2.52
	2007	92	2.26	1.82 -	2.77
	2008	73	1.80	1.41 -	2.26
	2009	72	1.79	1.40 -	2.26

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence	
Anotia or microtia	2000	100	2.75	2.24 -	3.35
	2001	105	2.88	2.33 -	3.43
	2002	106	2.85	2.30 -	3.39
	2003	102	2.70	2.18 -	3.23
	2004	130	3.41	2.82 -	3.99
	2005	114	2.96	2.41 -	3.50
	2006	134	3.36	2.79 -	3.92
	2007	124	3.04	2.51 -	3.58
	2008	164	4.05	3.43 -	4.67
	2009	134	3.34	2.77 -	3.90
Cardiac and Circulatory					
Common truncus	2000	27	0.74	0.49 -	1.08
	2001	26	0.71	0.47 -	1.04
	2002	29	0.78	0.52 -	1.12
	2003	28	0.74	0.49 -	1.07
	2004	31	0.81	0.55 -	1.15
	2005	24	0.62	0.40 -	0.93
	2006	28	0.70	0.47 -	1.01
	2007	35	0.86	0.60 -	1.19
	2008	48	1.18	0.87 -	1.57
	2009	32	0.80	0.55 -	1.12
Transposition of the great vessels	2000	163	4.49	3.80 -	5.18
	2001	193	5.29	4.54 -	6.03
	2002	164	4.40	3.73 -	5.08
	2003	197	5.22	4.49 -	5.95
	2004	182	4.77	4.08 -	5.46
	2005	176	4.57	3.89 -	5.24
	2006	209	5.23	4.52 -	5.94
	2007	202	4.96	4.27 -	5.64
	2008	222	5.48	4.76 -	6.20
	2009	221	5.50	4.78 -	6.23
Tetralogy of Fallot	2000	104	2.86	2.31 -	3.41
	2001	127	3.48	2.87 -	4.08
	2002	132	3.54	2.94 -	4.15
	2003	144	3.82	3.19 -	4.44
	2004	110	2.88	2.34 -	3.42
	2005	152	3.94	3.32 -	4.57
	2006	152	3.81	3.20 -	4.41
	2007	135	3.31	2.75 -	3.87
	2008	180	4.44	3.79 -	5.09
	2009	149	3.71	3.11 -	4.31

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence
Ventricular septal defect	2000	1555	42.80	40.67 - 44.93
	2001	1669	45.71	43.52 - 47.91
	2002	1824	48.98	46.74 - 51.23
	2003	1848	48.97	46.74 - 51.20
	2004	1969	51.62	49.34 - 53.90
	2005	2205	57.19	54.81 - 59.58
	2006	2277	57.02	54.68 - 59.37
	2007	2485	60.99	58.59 - 63.39
	2008	2621	64.68	62.20 - 67.15
	2009	2684	66.83	64.30 - 69.36
Atrial septal defect	2000	1426	39.25	37.21 - 41.29
	2001	1523	41.72	39.62 - 43.81
	2002	1620	43.51	41.39 - 45.62
	2003	1809	47.94	45.73 - 50.15
	2004	2136	56.00	53.62 - 58.37
	2005	2501	64.87	62.33 - 67.41
	2006	2469	61.83	59.39 - 64.27
	2007	2463	60.45	58.06 - 62.84
	2008	2808	69.29	66.73 - 71.85
	2009	2676	66.63	64.11 - 69.16
Atrioventricular septal defect (endocardial cushion defect)	2000	153	4.21	3.54 - 4.88
	2001	159	4.36	3.68 - 5.03
	2002	151	4.06	3.41 - 4.70
	2003	146	3.87	3.24 - 4.50
	2004	152	3.98	3.35 - 4.62
	2005	156	4.05	3.41 - 4.68
	2006	181	4.53	3.87 - 5.19
	2007	157	3.85	3.25 - 4.46
	2008	199	4.91	4.23 - 5.59
	2009	182	4.53	3.87 - 5.19
Pulmonary valve atresia or stenosis	2000	239	6.58	5.74 - 7.41
	2001	274	7.50	6.62 - 8.39
	2002	251	6.74	5.91 - 7.57
	2003	318	8.43	7.50 - 9.35
	2004	344	9.02	8.07 - 9.97
	2005	349	9.05	8.10 - 10.00
	2006	347	8.69	7.78 - 9.60
	2007	405	9.94	8.97 - 10.91
	2008	469	11.57	10.53 - 12.62
	2009	408	10.16	9.17 - 11.15

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence
Tricuspid valve atresia or stenosis	2000	42	1.16	0.83 - 1.56
	2001	57	1.56	1.18 - 2.02
	2002	56	1.50	1.14 - 1.95
	2003	67	1.78	1.38 - 2.25
	2004	70	1.84	1.43 - 2.32
	2005	60	1.56	1.19 - 2.00
	2006	79	1.98	1.57 - 2.47
	2007	77	1.89	1.49 - 2.36
	2008	77	1.90	1.50 - 2.37
	2009	73	1.82	1.42 - 2.29
Ebstein anomaly	2000	27	0.74	0.49 - 1.08
	2001	19	0.52	0.31 - 0.81
	2002	23	0.62	0.39 - 0.93
	2003	31	0.82	0.56 - 1.17
	2004	41	1.07	0.77 - 1.46
	2005	23	0.60	0.38 - 0.90
	2006	25	0.63	0.41 - 0.92
	2007	24	0.59	0.38 - 0.88
	2008	29	0.72	0.48 - 1.03
	2009	23	0.57	0.36 - 0.86
Aortic valve stenosis	2000	76	2.09	1.65 - 2.62
	2001	84	2.30	1.84 - 2.85
	2002	90	2.42	1.94 - 2.97
	2003	95	2.52	2.04 - 3.08
	2004	99	2.60	2.11 - 3.16
	2005	76	1.97	1.55 - 2.47
	2006	92	2.30	1.86 - 2.83
	2007	89	2.18	1.75 - 2.69
	2008	108	2.67	2.16 - 3.17
	2009	95	2.37	1.91 - 2.89
Hypoplastic left heart syndrome	2000	80	2.20	1.75 - 2.74
	2001	67	1.84	1.42 - 2.33
	2002	90	2.42	1.94 - 2.97
	2003	74	1.96	1.54 - 2.46
	2004	81	2.12	1.69 - 2.64
	2005	77	2.00	1.58 - 2.50
	2006	87	2.18	1.75 - 2.69
	2007	80	1.96	1.56 - 2.44
	2008	89	2.20	1.76 - 2.70
	2009	85	2.12	1.69 - 2.62

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence	
Patent ductus arteriosus	2000	1497	41.20	39.12 -	43.29
	2001	1608	44.04	41.89 -	46.20
	2002	1639	44.02	41.88 -	46.15
	2003	1800	47.70	45.49 -	49.90
	2004	2044	53.59	51.26 -	55.91
	2005	2010	52.14	49.86 -	54.41
	2006	2177	54.52	52.23 -	56.81
	2007	2133	52.35	50.13 -	54.57
	2008	2506	61.84	59.42 -	64.26
	2009	2625	65.36	62.86 -	67.86
Coarctation of the aorta	2000	147	4.05	3.39 -	4.70
	2001	155	4.25	3.58 -	4.91
	2002	194	5.21	4.48 -	5.94
	2003	199	5.27	4.54 -	6.01
	2004	181	4.75	4.05 -	5.44
	2005	204	5.29	4.57 -	6.02
	2006	226	5.66	4.92 -	6.40
	2007	195	4.79	4.11 -	5.46
	2008	210	5.18	4.48 -	5.88
	2009	200	4.98	4.29 -	5.67
Respiratory					
Choanal atresia or stenosis	2000	48	1.32	0.97 -	1.75
	2001	42	1.15	0.83 -	1.55
	2002	45	1.21	0.88 -	1.62
	2003	38	1.01	0.71 -	1.38
	2004	32	0.84	0.57 -	1.18
	2005	43	1.12	0.81 -	1.50
	2006	59	1.48	1.12 -	1.91
	2007	44	1.08	0.78 -	1.45
	2008	54	1.33	1.00 -	1.74
	2009	55	1.37	1.03 -	1.78
Agenesis, aplasia, or hypoplasia of the lung	2000	157	4.32	3.65 -	5.00
	2001	157	4.30	3.63 -	4.97
	2002	136	3.65	3.04 -	4.27
	2003	122	3.23	2.66 -	3.81
	2004	99	2.60	2.11 -	3.16
	2005	110	2.85	2.32 -	3.39
	2006	110	2.75	2.24 -	3.27
	2007	104	2.55	2.06 -	3.04
	2008	93	2.29	1.85 -	2.81
	2009	114	2.84	2.32 -	3.36

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries
 Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence	
Oral Clefts					
Cleft palate alone (without cleft lip)	2000	228	6.28	5.46 -	7.09
	2001	216	5.92	5.13 -	6.71
	2002	197	5.29	4.55 -	6.03
	2003	203	5.38	4.64 -	6.12
	2004	210	5.51	4.76 -	6.25
	2005	254	6.59	5.78 -	7.40
	2006	222	5.56	4.83 -	6.29
	2007	239	5.87	5.12 -	6.61
	2008	264	6.51	5.73 -	7.30
	2009	250	6.23	5.45 -	7.00
Cleft lip with or without cleft palate	2000	401	11.04	9.96 -	12.12
	2001	401	10.98	9.91 -	12.06
	2002	389	10.45	9.41 -	11.48
	2003	390	10.33	9.31 -	11.36
	2004	452	11.85	10.76 -	12.94
	2005	430	11.15	10.10 -	12.21
	2006	447	11.19	10.16 -	12.23
	2007	434	10.65	9.65 -	11.65
	2008	442	10.91	9.89 -	11.92
	2009	377	9.39	8.44 -	10.34
Gastrointestinal					
Tracheoesophageal fistula/esophageal atresia	2000	72	1.98	1.55 -	2.50
	2001	72	1.97	1.54 -	2.48
	2002	86	2.31	1.85 -	2.85
	2003	78	2.07	1.63 -	2.58
	2004	66	1.73	1.34 -	2.20
	2005	79	2.05	1.62 -	2.55
	2006	81	2.03	1.61 -	2.52
	2007	88	2.16	1.73 -	2.66
	2008	79	1.95	1.54 -	2.43
	2009	89	2.22	1.78 -	2.73
Pyloric stenosis	2000	747	20.56	19.09 -	22.03
	2001	719	19.69	18.25 -	21.13
	2002	645	17.32	15.98 -	18.66
	2003	584	15.48	14.22 -	16.73
	2004	776	20.34	18.91 -	21.78
	2005	691	17.92	16.59 -	19.26
	2006	980	24.54	23.01 -	26.08
	2007	918	22.53	21.07 -	23.99
	2008	715	17.64	16.35 -	18.94
	2009	624	15.54	14.32 -	16.76

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence
Stenosis or atresia of the small intestine	2000	112	3.08	2.51 - 3.65
	2001	130	3.56	2.95 - 4.17
	2002	111	2.98	2.43 - 3.54
	2003	96	2.54	2.06 - 3.11
	2004	126	3.30	2.73 - 3.88
	2005	114	2.96	2.41 - 3.50
	2006	130	3.26	2.70 - 3.82
	2007	151	3.71	3.11 - 4.30
	2008	147	3.63	3.04 - 4.21
	2009	138	3.44	2.86 - 4.01
Stenosis or atresia of large intestine, rectum, or anal canal	2000	163	4.49	3.80 - 5.18
	2001	190	5.20	4.46 - 5.94
	2002	201	5.40	4.65 - 6.14
	2003	213	5.64	4.89 - 6.40
	2004	192	5.03	4.32 - 5.75
	2005	214	5.55	4.81 - 6.29
	2006	210	5.26	4.55 - 5.97
	2007	213	5.23	4.53 - 5.93
	2008	220	5.43	4.71 - 6.15
	2009	218	5.43	4.71 - 6.15
Hirschsprung disease	2000	47	1.29	0.95 - 1.72
	2001	43	1.18	0.85 - 1.59
	2002	44	1.18	0.86 - 1.59
	2003	44	1.17	0.85 - 1.57
	2004	52	1.36	1.02 - 1.79
	2005	57	1.48	1.12 - 1.92
	2006	48	1.20	0.89 - 1.59
	2007	67	1.64	1.27 - 2.09
	2008	63	1.55	1.19 - 1.99
	2009	50	1.25	0.92 - 1.64
Biliary atresia	2000	28	0.77	0.51 - 1.11
	2001	24	0.66	0.42 - 0.98
	2002	24	0.64	0.41 - 0.96
	2003	31	0.82	0.56 - 1.17
	2004	28	0.73	0.49 - 1.06
	2005	26	0.67	0.44 - 0.99
	2006	33	0.83	0.57 - 1.16
	2007	30	0.74	0.50 - 1.05
	2008	26	0.64	0.42 - 0.94
	2009	32	0.80	0.55 - 1.12

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence [†]	95% Confidence Interval for Prevalence
Genitourinary				
Hypospadias (cases and prevalence among males)	2000	1030	55.50	52.11 - 58.89
	2001	981	52.52	49.24 - 55.81
	2002	986	51.85	48.61 - 55.09
	2003	1007	52.29	49.06 - 55.52
	2004	1042	53.43	50.19 - 56.67
	2005	1133	57.37	54.03 - 60.71
	2006	1103	54.06	50.87 - 57.25
	2007	1151	55.28	52.08 - 58.47
	2008	1211	58.36	55.07 - 61.65
	2009	1259	61.45	58.06 - 64.85
Epispadias	2000	32	0.88	0.60 - 1.24
	2001	29	0.79	0.53 - 1.14
	2002	22	0.59	0.37 - 0.89
	2003	35	0.93	0.65 - 1.29
	2004	30	0.79	0.53 - 1.12
	2005	26	0.67	0.44 - 0.99
	2006	41	1.03	0.74 - 1.39
	2007	37	0.91	0.64 - 1.25
	2008	40	0.99	0.71 - 1.34
	2009	39	0.97	0.69 - 1.33
Renal agenesis or dysgenesis	2000	177	4.87	4.15 - 5.59
	2001	193	5.29	4.54 - 6.03
	2002	181	4.86	4.15 - 5.57
	2003	222	5.88	5.11 - 6.66
	2004	212	5.56	4.81 - 6.31
	2005	189	4.90	4.20 - 5.60
	2006	240	6.01	5.25 - 6.77
	2007	247	6.06	5.31 - 6.82
	2008	241	5.95	5.20 - 6.70
	2009	253	6.30	5.52 - 7.08
Bladder exstrophy	2000	11	0.30	0.15 - 0.54
	2001	5	0.14	0.04 - 0.32
	2002	12	0.32	0.17 - 0.56
	2003	11	0.29	0.15 - 0.52
	2004	10	0.26	0.13 - 0.48
	2005	3	0.08	0.02 - 0.23
	2006	6	0.15	0.06 - 0.33
	2007	7	0.17	0.07 - 0.35
	2008	8	0.20	0.09 - 0.39
	2009	5	0.12	0.04 - 0.29

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence [†]	95% Confidence Interval for Prevalence	
Musculoskeletal					
Congenital hip dislocation	2000	171	4.71	4.00 -	5.41
	2001	152	4.16	3.50 -	4.83
	2002	138	3.71	3.09 -	4.32
	2003	173	4.58	3.90 -	5.27
	2004	174	4.56	3.88 -	5.24
	2005	179	4.64	3.96 -	5.32
	2006	186	4.66	3.99 -	5.33
	2007	195	4.79	4.11 -	5.46
	2008	162	4.00	3.38 -	4.61
	2009	156	3.88	3.27 -	4.49
Talipes equinovarus / clubfoot	2000	519	14.28	13.06 -	15.51
	2001	536	14.68	13.44 -	15.92
	2002	572	15.36	14.10 -	16.62
	2003	574	15.21	13.97 -	16.45
	2004	583	15.28	14.04 -	16.52
	2005	553	14.34	13.15 -	15.54
	2006	580	14.53	13.34 -	15.71
	2007	626	15.36	14.16 -	16.57
	2008	621	15.32	14.12 -	16.53
	2009	631	15.71	14.49 -	16.94
Reduction defects of the upper limbs	2000	157	4.32	3.65 -	5.00
	2001	149	4.08	3.43 -	4.74
	2002	148	3.97	3.33 -	4.61
	2003	151	4.00	3.36 -	4.64
	2004	149	3.91	3.28 -	4.53
	2005	156	4.05	3.41 -	4.68
	2006	165	4.13	3.50 -	4.76
	2007	167	4.10	3.48 -	4.72
	2008	168	4.15	3.52 -	4.77
	2009	164	4.08	3.46 -	4.71
Reduction defects of the lower limbs	2000	55	1.51	1.14 -	1.97
	2001	79	2.16	1.71 -	2.70
	2002	63	1.69	1.30 -	2.16
	2003	75	1.99	1.56 -	2.49
	2004	64	1.68	1.29 -	2.14
	2005	83	2.15	1.71 -	2.67
	2006	73	1.83	1.43 -	2.30
	2007	89	2.18	1.75 -	2.69
	2008	83	2.05	1.63 -	2.54
	2009	70	1.74	1.36 -	2.20

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence
Craniosynostosis	2000	163	4.49	3.80 - 5.18
	2001	139	3.81	3.17 - 4.44
	2002	173	4.65	3.95 - 5.34
	2003	168	4.45	3.78 - 5.13
	2004	172	4.51	3.84 - 5.18
	2005	159	4.12	3.48 - 4.77
	2006	180	4.51	3.85 - 5.17
	2007	217	5.33	4.62 - 6.03
	2008	213	5.26	4.55 - 5.96
	2009	201	5.00	4.31 - 5.70
Achondroplasia	2000	10	0.28	0.13 - 0.51
	2001	13	0.36	0.19 - 0.61
	2002	10	0.27	0.13 - 0.49
	2003	11	0.29	0.15 - 0.52
	2004	9	0.24	0.11 - 0.45
	2005	10	0.26	0.12 - 0.48
	2006	9	0.23	0.10 - 0.43
	2007	19	0.47	0.28 - 0.73
	2008	12	0.30	0.15 - 0.52
	2009	15	0.37	0.21 - 0.62
Diaphragmatic hernia	2000	98	2.70	2.19 - 3.29
	2001	83	2.27	1.81 - 2.82
	2002	96	2.58	2.09 - 3.15
	2003	108	2.86	2.32 - 3.40
	2004	100	2.62	2.13 - 3.19
	2005	98	2.54	2.06 - 3.10
	2006	116	2.91	2.38 - 3.43
	2007	128	3.14	2.60 - 3.69
	2008	115	2.84	2.32 - 3.36
	2009	107	2.66	2.16 - 3.17
Omphalocele	2000	83	2.28	1.82 - 2.83
	2001	85	2.33	1.86 - 2.88
	2002	77	2.07	1.63 - 2.58
	2003	73	1.93	1.52 - 2.43
	2004	83	2.18	1.73 - 2.70
	2005	73	1.89	1.48 - 2.38
	2006	86	2.15	1.72 - 2.66
	2007	80	1.96	1.56 - 2.44
	2008	79	1.95	1.54 - 2.43
	2009	89	2.22	1.78 - 2.73

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries

Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence	
Gastroschisis	2000	143	3.94	3.29 -	4.58
	2001	148	4.05	3.40 -	4.71
	2002	149	4.00	3.36 -	4.64
	2003	173	4.58	3.90 -	5.27
	2004	165	4.33	3.67 -	4.99
	2005	208	5.40	4.66 -	6.13
	2006	204	5.11	4.41 -	5.81
	2007	257	6.31	5.54 -	7.08
	2008	250	6.17	5.40 -	6.93
	2009	242	6.03	5.27 -	6.79
Chromosomal					
Trisomy 21 (Down syndrome)	2000	476	13.10	11.92 -	14.28
	2001	494	13.53	12.34 -	14.72
	2002	454	12.19	11.07 -	13.31
	2003	457	12.11	11.00 -	13.22
	2004	513	13.45	12.29 -	14.61
	2005	482	12.50	11.39 -	13.62
	2006	559	14.00	12.84 -	15.16
	2007	552	13.55	12.42 -	14.68
	2008	544	13.42	12.30 -	14.55
	2009	550	13.70	12.55 -	14.84
Trisomy 13 (Patau syndrome)	2000	51	1.40	1.05 -	1.85
	2001	42	1.15	0.83 -	1.55
	2002	43	1.15	0.84 -	1.56
	2003	42	1.11	0.80 -	1.50
	2004	47	1.23	0.91 -	1.64
	2005	34	0.88	0.61 -	1.23
	2006	42	1.05	0.76 -	1.42
	2007	51	1.25	0.93 -	1.65
	2008	45	1.11	0.81 -	1.49
	2009	53	1.32	0.99 -	1.73
Trisomy 18 (Edwards syndrome)	2000	78	2.15	1.70 -	2.68
	2001	81	2.22	1.76 -	2.76
	2002	82	2.20	1.75 -	2.73
	2003	80	2.12	1.68 -	2.64
	2004	96	2.52	2.04 -	3.07
	2005	96	2.49	2.02 -	3.04
	2006	110	2.75	2.24 -	3.27
	2007	100	2.45	2.00 -	2.99
	2008	120	2.96	2.43 -	3.49
	2009	115	2.86	2.34 -	3.39

Texas Birth Defects Registry, Report of Defects Among 2000-2009 Deliveries
 Table 2. Prevalence of Selected Birth Defects by Year, Texas, 2000-2009

Defect	Year	Cases	Prevalence[†]	95% Confidence Interval for Prevalence
Infants and fetuses with any monitored birth defect	2000	13034	358.74	352.58 - 364.90
	2001	12975	355.39	349.27 - 361.51
	2002	13773	369.88	363.70 - 376.05
	2003	14399	381.56	375.33 - 387.79
	2004	15814	414.59	408.12 - 421.05
	2005	16693	432.98	426.41 - 439.55
	2006	17410	436.00	429.53 - 442.48
	2007	18119	444.69	438.21 - 451.16
	2008	18964	467.97	461.31 - 474.63
	2009	19697	490.46	483.61 - 497.31

[†] cases per 10,000 live births