

	Population	433,082	1,674,050	2,010,968	2,139,780	2,202,367	4,231,136	4,107,353	3,780,922	3,537,301	5,249,520	29,366,479
DISEASE	<1	1-4	5-9	10-14	15-19	20-29	30-39	40-49	50-59	60+	TOTAL	
TRICHURIASIS	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
TYPHOID FEVER	0.0	0.2	0.0	0.2	0.0	0.2	0.0	0.2	0.1	0.0	0.1	
TYPHUS, MURINE	0.2	1.3	3.2	3.9	3.5	1.9	2.6	2.4	2.7	2.3	2.5	
VIBRIO PARAHAEMOLYTICUS	0.0	0.1	0.0	0.1	0.1	0.2	0.2	0.3	0.1	0.1	0.1	
VIBRIO VULNIFICUS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.1	
VIBRIO, OTHER/UNSPECIFIED	3.9	0.8	0.6	0.2	0.3	0.3	0.4	0.5	0.6	1.6	0.7	
VISA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
WEST NILE FEVER	0.0	0.0	0.0	0.0	0.2	0.1	0.1	0.1	0.3	0.2	0.1	
WEST NILE NEUROINVASIVE DISEASE	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.6	0.3	1.3	0.4	
YERSINIOSIS	0.5	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.3	0.1	
ZIKA VIRUS DISEASE	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	

¹ Population data is projected population accessed 4/18/18 from the Department of State Health Services, Center for Health Statistics <http://www.dshs.state.tx.us/chs/popdat/popup.shtm>.

² Infant botulism and congenital rubella syndrome rates are calculated using the Texas populations under 1 year of age.

³ Perinatal hepatitis B cases are defined as infants >1 months through 24 months of age. However, the rate calculation for those <1 year does not exclude those <1 month of age. The rate calculation for the 1-4 year age group cell includes only the 1 year old population.

⁴ Influenza-associated pediatric mortality rates were calculated for the Texas populations <18 years of age including those <1, 1-4, 5-9, 10-14, and for the 15-19 age group only those 15-17.